arXiv:1303.2839v1 [math-ph] 12 Mar 2013

# A Note Basis Properties for Fractional Hydrogen Atom Equation

E. BAS[a], F. METIN[b]

[a,b]Department of Mathematics, Faculty of Science, Firat University, Elazig, 23119, Turkey

Reprint requests to E. B.; E-mail: erdalmat@yahoo.com.

In this paper, spectral analysis of fractional Sturm Liouville problem defined on $(0,1]$, having the singularity of type $-\frac{2}{r}+\frac{l(l+1)}{r^2}$ at zero and research the fundamental properties of the eigenfunctions and eigenvalues for the operator. We show that the eigenvalues and eigenfunctions of the problem are real and orthogonal, respectively. Furthermore,we give some important theorems and lemmas for fractional hydrogen atom equation.

Keywords: Sturm-Liouville, Fractional, Hydrogen atom, Singular, Spectral.

## 1. Introduction

Sturm-Liouville problem was first developed in a number of papers that were published by these authors in 1836 and 1837. (Charles-François Sturm (1803-1855), Professor of Mechanics at the Sorbonne, had been interested, since about 1833, in the problem of heat flow in bars, so was well aware of eigenvalue-type problems. He worked closely with his friend Joseph Liouville (1809-1882), Professor of Mathematics at the Collège de France, on the general properties of second order differential equations. Liouville also made many contributions to the general field of analysis.) *[1]*. Mathematical techniques could be developed into a more suitable and significant course by presenting them within the more general Sturm-Liouville theory in $L_2$. The Sturm-Liouville Problems are important in many areas of science, engineering and mathematics. It is known that the spectral characteristics are spectra, spectral functions, scattering data, norming constants, etc. According to the theory *a* linear second-order differential operator which is self-adjoint has an orthogonal sequence of eigenfunctions in $L_2$. Spectral properties of Sturm-Liouville operators are often derived, directly or indirectly, as a consequence of an established link between large distance asymptotic behavior of solutions of the associated differential equation and spectral properties of the corresponding differential operator. Sturm-Liouville problems are divided into regular and singular type. Differential equations such as Bessel, hydrogen atom, Hermitte, Jakobi, and Legendre equations can be transformed into Sturm Liouville equations. There are many studies on these issues, *[1-5]*. We also discuss the radial part of Schrödinger's equation for the hydrogen atom.

Fractional calculus is 'the theory of derivatives and integrals of any arbitrary real or complex order, which unify and generalize the notions of integer-order differentiation and $n-$fold integration' *[6-13]*. In recent years, the concept of fractional calculus, originated from Leibniz, has achieved increasing interest during last two decades. In particular, on the last decade have scientific papers concerning fractional quantum mechanics. It has been proved that many systems in different fields of science and engineering can be modeled more accurately using fractional derivatives *[14,15]*. We note that ordinary derivatives in a traditional Sturm-Liouville problem are replaced with fractional derivatives and the resulting problems are solved using some numerical methods *[16-20]*. Furthermore, Klimek and Agrawal *[21]* define a fractional Sturm-Liouville operator, introduce a regular fractional Sturm-Liouville probelem and investigate the properties of the eigenfunctions and the eigenvalues of the operator. In this paper, our purpose is to introduce regular fractional singular Sturm-Liouville problem and prove spectral properties of spectral data for the operator.

Let's give the main problem and necessary datas, as follows:

## 2. Preliminaries

Firstly, we consider singular Sturm-Liouville equation

$$\frac{d^2R}{dr^2}+\frac{a}{r}\frac{dR}{dr}-\frac{\ell(\ell+1)}{r^2}R+\left(E+\frac{a}{r}\right)R=0 \qquad (0<r<\infty).$$

In quantum mechanics the study of the energy levels of the hydrogen atom leads to this equation [5]. Where $R$ is the distance of the mass center to the origin, $\ell$ is positive integer, $a$ is real number $E$ is energy constant and $r$ is the distance between the nucleus with the electron.

Substitution $R=y/r$ reduces above equation to the form

$$\frac{d^2y}{dr^2}+\left\{E+\frac{a}{r}-\frac{\ell\left(\ell+1\right)}{r^2}\right\}y=0.$$

In most cases we took the reference potential to be

$$V\left(r\right)=\frac{a}{r}-\frac{\ell\left(\ell+1\right)}{r^2}$$

when $V\left(r\right)$ is defined by last differential equation, it contains a centripetal and Colomb part, the usual singularities of the nuclear problem.

**Definition 1.** Let $\operatorname{Re}\left(\alpha\right)>0$. The left-sided and respectively right-sided Riemann-Liouville integrals of order $\alpha$ are given by the formulas

$$\left(I_{a,+}^{\alpha}f\right)\left(r\right)=\frac{1}{\Gamma\left(\alpha\right)}\int\limits_a^r\left(r-s\right)^{\alpha-1}f\left(s\right)ds,\quad r>a, \tag{1}$$

$$\left(I_{b,-}^{\alpha}f\right)\left(r\right)=\frac{1}{\Gamma\left(\alpha\right)}\int\limits_r^b\left(s-r\right)^{\alpha-1}f\left(s\right)ds,\quad r<b \tag{2}$$

where $\Gamma$ denotes the gamma function.

**Definition 2.** Let $\operatorname{Re}\left(\alpha\right)>0$. The left-sided and respectively right-sided Riemann-Liouville derivatives of order $\alpha$ are defined as

$$\left(D_{a,+}^{\alpha}f\right)\left(r\right)=D\left(I_{a,+}^{1-\alpha}f\right)\left(r\right)\qquad r>a, \tag{3}$$

$$\left(D_{b,-}^{\alpha}f\right)\left(r\right)=-D\left(I_{b,-}^{1-\alpha}f\right)\left(r\right)\quad r<b, \tag{4}$$

analogous formulas yield the left- and right-sided Caputo derivatives of order $\alpha$:

$$\left({}^{C}D_{a,+}^{\alpha}f\right)\left(r\right)=\left(I_{a,+}^{1-\alpha}Df\right)\left(r\right)\qquad r>a, \tag{5}$$

$$\left({}^{C}D_{b,-}^{\alpha}f\right)\left(r\right)=\left(I_{b,-}^{1-\alpha}\left(-D\right)f\right)\left(r\right)\quad r<b. \tag{6}$$

**Definition 3.***[14]* The general function ${}_p\Psi_q\left(z\right)$ is defined for $z\in\mathbb{C}$, complex $a_l,b_j\in\mathbb{C}$, and real $\alpha_i,\beta_j\in\mathbb{R}\left(i=1,...,p;j=1,...q\right)$ by the series

$$\begin{aligned}{}_p\Psi_q\left(z\right)&={}_p\ \Psi_q\left[\begin{array}{c}\left(a_1,\alpha_1\right)_{1,p}\\\left(b_1,\beta_1\right)_{1,q}\end{array}\middle|z\right]\\&=\sum_{k=0}^{\infty}\frac{\prod\limits_{i=1}^{p}\Gamma\left(a_i+\alpha_lk\right)}{\prod\limits_{j=1}^{q}\Gamma\left(b_j+\beta_jk\right)}\frac{z^k}{k!}\end{aligned} \tag{7}$$

where $z,a_i,b_j\in\mathbb{C},\alpha_i,\beta_j\in\mathbb{R}\left(i=1,...,p;j=1,...q\right)$. This general Wright function was investigated by Fox who presented its asymptotic expansion for large values of the argument $z$ under the condition

$$\sum_{j=1}^{q}\beta_j-\sum_{i=1}^{p}\alpha_i>1.$$

If these conditions are satisfied, the series in (7) is convergent for any $z\in\mathbb{C}$.

**Theorem 4.** *[14]* Let $a_l,b_j\in\mathbb{C},\alpha_l,\beta_j\in\mathbb{R}\ \left(i=1,...,p\right)$, $\left(j=1,...q\right)$ and let

$$\Delta=\sum_{j=1}^{q}\beta_j-\sum_{l=1}^{p}\alpha_i,$$

$$\delta=\prod_{i=1}^{p}\left|\alpha_i\right|^{-\alpha_l}\prod_{j=1}^{q}\left|\beta_j\right|^{\beta_j},$$

$$\mu=\sum_{j=1}^{q}b_j-\sum_{l=1}^{p}a_i+\frac{p-q}{2}$$

**I)** If $\Delta>-1$, then the series in (7) is absolutely convergent for all $z\in\mathbb{C}$.

**II)** If $\Delta=-1$, then the series in (7) is absolutely convergent for $\left|z\right|<\delta$ and for $\left|z\right|=\delta$ and $\Re\left(\mu\right)>1/2$.

**Definition 5.***[22]* Let $\left(X,d\right)$ be a non-empty complete metric space. Let $T:X\to X$ be a contraction mapping on $r$, i.e.. there is a nonnegative real number $q<1$ such that

$$d\left(T\left(r\right),T\left(y\right)\right)\leq\sigma d\left(r,y\right)$$

for all $r$, $y$ in $X$. Then the map $T$ admits one and only one fixed-point $r^*$ in $X$. (this means $T(r^*)=r^*$). Furthermore, this fixed point can be found as follows start with an arbitrary element $r_0$ in $r$ and define an iterative sequence by $r_n=T(r_{n-1})$ for $n=1,2,3,...$ This sequence converges, and its limit is $r^*$. The following inequality describes the speed of convergence:

$$d\left(r^*,r_n\right)\leq\frac{q_n}{1-q}d\left(r_1,r_0\right).$$

Equivalently,

$$d\left(r^*,r_{n+1}\right)\leq\frac{\sigma}{1-\sigma}d\left(r_{n+1},r_n\right)$$

and

$$d\left(r^*,r_{n+1}\right)\leq\sigma d\left(r^*,r_n\right).$$

Any such value of $\sigma$ is called a Lipschitz constant for $T$, and the smallest one is sometimes called "the best Lipschitz constant" of $T$.

**Property 6.** The fractional differential operators defined in $(3-6)$ satisfy the following identities:

$$\int\limits_a^b f\left(r\right)D_{b,-}^{\alpha}g\left(r\right)dr=\int\limits_a^b g\left(r\right)^{C}D_{a,+}^{\alpha}f\left(r\right)dr-f\left(r\right)I_{b,-}^{1-\alpha}g\left(r\right)\Big|_a^b$$

or

$$\int_a^b f(r)\, D^{\alpha}_{b,-} g(r)^{C} D^{\alpha}_{a,+} k(r)\, dr$$

$$= \int_a^b g(r)^{C} D^{\alpha}_{a,+} f(r)^{C} D^{\alpha}_{a,+} k(r)\, dr \tag{8}$$

$$- \; f(r)\, I^{1-\alpha}_{b,-} g(r)^{C} D^{\alpha}_{a,+} k(r) \;\Big|\; {}^{b}_{a},$$

$$\int_a^b f(r)\, D^{\alpha}_{a,+} g(r)\, dr$$

$$= \int_a^b g(r)^{C} D^{\alpha}_{b,-} f(r)\, dr + f(r)\, I^{1-\alpha}_{a,+} g(r)\, |^{b}_{a}\,. \tag{9}$$

**Property 7.** Assume $\alpha \in (0,1)$, $\beta > \alpha$ and $f \in C[a,b]$. Then the following relations

$$\left.\begin{array}{c} D^{\alpha}_{a,+} I^{\alpha}_{a,+} f(r) = f(r) \\ D^{\alpha}_{b,-} I^{\alpha}_{b,-} f(r) = f(r) \\ D^{\alpha}_{a,+} I^{\beta}_{a,+} f(r) = I^{\beta-\alpha}_{a,+} f(r) \\ D^{\alpha}_{b,-} I^{\beta}_{b,-} f(r) = I^{\beta-\alpha}_{b,-} f(r) \\ {}^{C}D^{\alpha}_{a,+} I^{\alpha}_{a,+} f(r) = f(r) \\ {}^{C}D^{\alpha}_{b,-} I^{\alpha}_{b,-} f(r) = f(r) \end{array}\right\} \tag{10}$$

hold for any $r \in [a,b]$. Furthermore, the integral operators defined in $(1,2)$ satisfy the following semi-group properties,

$$I^{\alpha}_{a,+} I^{\beta}_{a,+} = I^{\alpha+\beta}_{a,+}, \qquad I^{\alpha}_{b,-} I^{\beta}_{b,-} = I^{\alpha+\beta}_{b,-}$$

Now, let's take up a fractional Sturm-Liouville problem for hydrogen atom equation

## 3. A Singular Fractional Sturm-Liouville Problem for Hydrogen Atom Equation

Let's denote a singular fractional Sturm-Liouville problem for hydrogen atom equation with the differential part containing the left-and right-sided derivatives. Let's use the form of the integration by parts formulas $(8,9)$ for this new approximation. Main properties of eigenfunctions and eigenvalues in the theory of classical Sturm-Liouville problems are related to the integration by parts formula for the first order derivatives. In the corresponding fractional version we note that both left and right derivatives appear and the essential pairs are the left Riemann-Liouville derivative with the right Caputo derivative and the right Riemann-Liouville derivative with the left Caputo one.

**Definition 8.** For $\alpha \in (0,1)$, Fractional hydrogen atom operator is written as

$$\mathcal{L}_{\alpha} = D^{\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+} + \left( -\frac{2}{r} + \frac{l(l+1)}{r^2} + q_0(r) \right) \tag{11}$$

consider the fractional hydrogen atom equation

$$\mathcal{L}_{\alpha} y_{\lambda}(r) + \lambda w_{\alpha}(r)\, y_{\lambda}(r) = 0 \tag{12}$$

where $p(r) \neq 0, w_{\alpha}(r) > 0 \; \forall r \in (0,1]$ and $p, q$ are real valued continuous functions in interval $(0,1]$. The boundary conditions for the operator $\mathcal{L}_{\alpha}$ are the following:

$$y_{\lambda}(0) = 0, \tag{13}$$

$$d_1 y(1) + d_2 I^{1-\alpha}_{1,-} p(1)^{C} D^{\alpha}_{0,+} y(1) = 0, \tag{14}$$

where $d_1^2 + d_2^2 \neq 0$. The fractional boundary-value problem (12)-(14) is fractional Sturm-Liouville problem for hydrogen atom operator.

**Theorem 9.** Fractional hydrogen operator is self-adjoint on $(0,1]$.

**Proof.** Let us consider the following equation

$$\langle \mathcal{L}_{\alpha}\varphi, \phi \rangle = \int_0^1 \mathcal{L}_{\alpha}\varphi\phi dr = \int_0^1 \phi(r) \Big[ D^{\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+} \varphi(r)$$

$$+ \left( -\frac{2}{r} + \frac{l(l+1)}{r^2} + q_0(r) \right) \varphi(r) \Big] dr$$

$$= \int_0^1 \phi(r)\, D^{\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+} \varphi(r)\, dr$$

$$+ \int_0^1 \left( -\frac{2}{r} + \frac{l(l+1)}{r^2} + q_0(r) \right) \varphi(r)\, \phi(r)\, dr.$$

Considering property (8) to the first integral in the last equation, we obtain the identity

$$
\begin{aligned}
\langle \mathcal{L}_\alpha \varphi, \phi \rangle &= \int_0^1 p\left(r\right) {}^{C}D^{\alpha}_{0,+}\phi\left(r\right) {}^{C}D^{\alpha}_{0,+}\varphi\left(r\right) dr \\
&-\phi\left(r\right) I^{1-\alpha}_{1,-} p\left(r\right) {}^{C}D^{\alpha}_{0,+}\varphi\left(r\right)\Big|_0^1 \\
&+\int_0^1 \left(-\frac{2}{r}+\frac{l\left(l+1\right)}{r^2}+q_0\left(r\right)\right)\varphi\left(r\right)\phi\left(r\right) dr \\
&= \int_0^1 p\left(r\right) {}^{C}D^{\alpha}_{0,+}\phi\left(r\right) {}^{C}D^{\alpha}_{0,+}\varphi\left(r\right) dr \\
&+\frac{d_1}{d_2}\varphi\left(1\right)\phi\left(1\right)+\frac{c_1}{c_2}\varphi\left(0\right)\phi\left(0\right) \\
&+\int_0^1 \left(-\frac{2}{r}+\frac{l\left(l+1\right)}{r^2}+q_0\left(r\right)\right)\varphi\left(r\right)\phi\left(r\right) dr. \quad (15)
\end{aligned}
$$

Similarly, we obtain

$$
\begin{aligned}
\langle \varphi, \mathcal{L}_\alpha \phi \rangle &= \int_0^1 p\left(r\right) {}^{C}D^{\alpha}_{0,+}\varphi\left(r\right) {}^{C}D^{\alpha}_{0,+}\phi\left(r\right) dr \\
&+\frac{d_1}{d_2}\varphi\left(1\right)\phi\left(1\right)+\frac{c_1}{c_2}\varphi\left(0\right)\phi\left(0\right) \\
&+\int_0^1 \left(-\frac{2}{r}+\frac{l\left(l+1\right)}{r^2}+q_0\left(r\right)\right)\phi\left(r\right)\varphi\left(r\right) dr. \quad (16)
\end{aligned}
$$

The right hand sides of the equations (15) and (16) are equal hence we may see that the left sides are equal that is

$$\langle \mathcal{L}_\alpha \varphi, \phi \rangle = \langle \varphi, \mathcal{L}_\alpha \phi \rangle .$$

Therefore, $\mathcal{L}_\alpha = \mathcal{L}^*_\alpha$. The proof is completed.

**Theorem 10.** The eigenvalues of fractional hydrogen atom operator $(12-14)$ are real.

**Proof:** Let us observe that following relation results from Property (8)

$$
\begin{aligned}
\int_0^1 f\left(r\right)\mathcal{L}_\alpha g\left(r\right) dr \; &= \int_0^1 p\left(r\right) {}^{C}D^{\alpha}_{0,+} f\left(r\right) {}^{C}D^{\alpha}_{0,+} g\left(r\right) dr \\
&-f\left(r\right) I^{1-\alpha}_{1,-} p\left(r\right) {}^{C}D^{\alpha}_{0,+} g\left(r\right)\Big|_0^1 \\
&+\int_0^1 \left(-\frac{2}{r}+\frac{l\left(l+1\right)}{r^2}+q_0\left(r\right)\right) g\left(r\right) f\left(r\right) dr. \qquad (17)
\end{aligned}
$$

Suppose that $\lambda$ is the eigenvalue for $(12-14)$ corresponding to eigenfunction $y$ the following equalities are satisfy $y$ and its complex conjugate $\bar{y}$

$$\mathcal{L}_\alpha y\left(r\right)+\lambda w_\alpha\left(r\right) y\left(r\right)=0 \qquad (18)$$

$$y_\lambda\left(0\right)=0 \qquad (19)$$

$$d_1 y\left(1\right)+d_2 I^{1-\alpha}_{1,-} p\left(1\right) {}^{C}D^{\alpha}_{0,+} y\left(1\right)=0 \qquad (20)$$

$$\mathcal{L}_\alpha \bar{y}\left(r\right)+\bar{\lambda} w_\alpha\left(r\right)\bar{y}\left(r\right)=0 \qquad (21)$$

$$\bar{y}\left(0\right)=0 \qquad (22)$$

$$d_1 \bar{y}\left(1\right)+d_2 I^{1-\alpha}_{1,-} p\left(1\right) {}^{C}D^{\alpha}_{0,+} \bar{y}\left(1\right)=0 \qquad (23)$$

where $d_1^2+d_2^2 \neq 0$. We multiply equation (18) by function $\bar{y}$ and (21) by function $y$ respectively and subtract:

$$\left(\lambda-\bar{\lambda}\right) w_\alpha\left(r\right) y\left(r\right)\bar{y}\left(r\right)=y\left(r\right)\mathcal{L}_\alpha \bar{y}\left(r\right)-\bar{y}\left(r\right)\mathcal{L}_\alpha y\left(r\right).$$

Now, we integrate over interval $(0,1]$ and applying relation (17) we note that the right-hand side of the integrated equality contains only boundary terms:

$$
\begin{aligned}
&\left(\lambda-\bar{\lambda}\right)\int_0^1 w_\alpha\left(r\right) y\left(r\right)\bar{y}\left(r\right) dr \\
&= \int_0^1 y\left(r\right)\mathcal{L}_\alpha \bar{y}\left(r\right) dr-\int_0^1 \bar{y}\left(r\right)\mathcal{L}_\alpha y\left(r\right) dr \\
&= \int_0^1 y\left(r\right)\Big[D^{\alpha}_{1,-} p\left(r\right) {}^{C}D^{\alpha}_{0,+}\bar{y}\left(r\right) \\
&+\left(-\frac{2}{r}+\frac{l\left(l+1\right)}{r^2}+q_0\left(r\right)\right)\bar{y}\left(r\right)\Big] dr \\
&-\int_0^1 \bar{y}\Big[D^{\alpha}_{1,-} p\left(r\right) {}^{C}D^{\alpha}_{0,+}\left(r\right) y\left(r\right) \\
&+\left(-\frac{2}{r}+\frac{l\left(l+1\right)}{r^2}+q_0\left(r\right)\right) y\left(r\right)\Big] dr \\
&= \int_0^1 p\left(r\right) {}^{C}D^{\alpha}_{0,+} y\left(r\right) {}^{C}D^{\alpha}_{0,+}\bar{y}\left(r\right) dr \\
&-y\left(r\right) I^{1-\alpha}_{1,-} p\left(r\right) {}^{C}D^{\alpha}_{0,+}\bar{y}\left(r\right)\Big|_0^1 \\
&+\int_0^1 \left(-\frac{2}{r}+\frac{l\left(l+1\right)}{r^2}+q_0\left(r\right)\right) y\left(r\right)\bar{y}\left(r\right) dr \\
&-\int_0^1 p\left(r\right) {}^{C}D^{\alpha}_{0,+}\bar{y}\left(r\right) {}^{C}D^{\alpha}_{0,+} y\left(r\right) dr
\end{aligned}
$$

or

$$+\bar{y}(r)\, I^{1-\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+} y(r)\Big|_{0}^{1}$$

$$-\int_{0}^{1}\left(-\frac{2}{r}+\frac{l(l+1)}{r^{2}}+q_{0}(r)\right) y(r)\,\bar{y}(r)\,dr$$

$$\left(\lambda-\bar{\lambda}\right)\int_{0}^{1} w_{\alpha}(r)\,|y(r)|^{2}\,dr$$

$$= -y(r)\, I^{1-\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+}\bar{y}(r)\Big|_{1}$$

$$+\, y(r)\, I^{1-\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+}\bar{y}(r)\Big|_{0}$$

$$+\bar{y}(r)\, I^{1-\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+} y(r)\Big|_{1}$$

$$-\,\bar{y}(r)\, I^{1-\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+} y(r)\Big|_{0}$$

by virtue of the boundary conditions $(19),(20),(22),(23)$, we find

$$\left(\lambda-\bar{\lambda}\right)\int_{0}^{1} w_{\alpha}(r)\,|y(r)|^{2}\,dr = 0$$

and because $y$ is a non-trivial solution and $w_{\alpha}(r) > 0$, it easily seen that $\lambda=\bar{\lambda}$.

**Theorem 11.** The eigenfunctions corresponding to distinct eigenvalues of fractional hydrogen atom operator $(12-14)$ are orthogonal weight function $w_{\alpha}$ on $(0,1]$ that is

$$\int_{0}^{1} w_{\alpha}(r)\, y_{\lambda_1}(r)\, y_{\lambda_2}(r)\,dr = 0 \qquad \lambda_1 \neq \lambda_2.$$

**Proof.** We have by assumptions fractional Sturm-Liouville for hydrogen atom operator fulfilled by two different eigenvalues $(\lambda_1,\lambda_2)$ and the respective eigenfunctions $(y_{\lambda_1},y_{\lambda_2})$, in that case

$$\mathcal{L}_{\alpha} y_{\lambda_1}(r) + \lambda_1 w_{\alpha}(r)\, y_{\lambda_1}(r) = 0 \tag{24}$$

$$y_{\lambda_1}(r) = 0 \tag{25}$$

$$d_1 y_{\lambda_1}(1) + d_2 I^{1-\alpha}_{1,-} p(1)^{C} D^{\alpha}_{0,+} y_{\lambda_1}(1) = 0 \tag{26}$$

$$\mathcal{L}_{\alpha} y_{\lambda_2}(r) + \lambda_2 w_{\alpha}(r)\, y_{\lambda_2}(r) = 0 \tag{27}$$

$$y_{\lambda_2}(r) = 0 \tag{28}$$

$$d_1 y_{\lambda_2}(1) + d_2 I^{1-\alpha}_{1,-} p(1)^{C} D^{\alpha}_{0,+} y_{\lambda_2}(1) = 0 \tag{29}$$

we multiply equation (24) by function $y_{\lambda_2}$ and (27) by function $y_{\lambda_1}$ respectively and subtract:

$$(\lambda_1-\lambda_2)\, w_{\alpha}(r)\, y_{\lambda_1} y_{\lambda_2} = y_{\lambda_1}\mathcal{L}_{\alpha} y_{\lambda_2} - y_{\lambda_2}\mathcal{L}_{\alpha} y_{\lambda_1}.$$

Integrating over interval $(0,1]$ and applying relation (17) we note that the right-hand side of the integrated equality contains only boundary terms:

$$(\lambda_1-\lambda_2)\int_{0}^{1} w_{\alpha}(r)\, y_{\lambda_1}(r)\, y_{\lambda_2}(r)\,dr$$

$$= \int_{0}^{1} y_{\lambda_1}(r)\,\mathcal{L}_{\alpha} y_{\lambda_2}(r)\,dr - \int_{0}^{1} y_{\lambda_2}(r)\,\mathcal{L}_{\alpha} y_{\lambda_1}(r)\,dr$$

$$= \int_{0}^{1} y_{\lambda_1}(r)\Big[D^{\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+} y_{\lambda_2}(r)$$

$$+\left(-\frac{2}{r}+\frac{l(l+1)}{r^{2}}+q_{0}(r)\right) y_{\lambda_2}(r)\Big]\,dr$$

$$-\int_{0}^{1} y_{\lambda_2}(r)\Big[D^{\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+}(r)\, y_{\lambda_1}(r)$$

$$+\left(-\frac{2}{r}+\frac{l(l+1)}{r^{2}}+q_{0}(r)\right) y_{\lambda_1}(r)\Big]\,dr$$

Using the boundary conditions $(25),(26),(28),(29)$, we find that

$$(\lambda_1-\lambda_2)\int_{0}^{1} w_{\alpha}(r)\, y_{\lambda_1}(r)\, y_{\lambda_2}(r)\,dr = 0,$$

where $\lambda_1 \neq \lambda_2$.

**Remark**. Let us now give certain auxiliary functions. Because we use the functions, the first of them is as follows

$$I^{\alpha}_{0,+}\frac{(1-r)^{\alpha-1}}{\Gamma(\alpha)} = (1-0)^{\alpha-1}(r-0)^{\alpha}$$

$$\cdot\, {}_1\Psi_2\left[\begin{array}{cc} (1,1) & \\ (\alpha,-1) & (\alpha+1,1)\end{array}\middle| -\frac{r-0}{1-0}\right]$$

where ${}_1\Psi_2$ is the Fox-Wright function *[14]*.

$${}_1\Psi_2\left[\begin{array}{cc} (a_1,\alpha_1) & \\ (b_1,\beta_1) & (b_2,\beta_2)\end{array}\middle| z\right]$$

$$= \sum_{k=0}^{\infty}\frac{\Gamma(a_1+\alpha_1 k)}{\Gamma(b_1+\beta_1 k)\,\Gamma(b_2+\beta_2 k)}\frac{z^{k}}{k!},$$

the properties of the function are determined by the parameters

$$\begin{aligned}\Delta &= \beta_1+\beta_2-\alpha_1 = -1\\ \delta &= |\alpha_1|^{-\alpha_1}\,|\beta_1|^{\beta_1}\,|\beta_2|^{\beta_2} = 1\\ \mu &= b_1+b_2-\alpha_1+\frac{1-2}{2} = 2\alpha-\frac{1}{2},\end{aligned} \tag{30}$$

considering Theorem 4, we note that this function is continuous in $(0,1]$ when order $\alpha > 1/2$, i.e. $\mu > 1/2$. It is discontinuous at end $r = 1$ for $0 < \alpha \le 1/2$. The explicitly calculated function allows to estimate the second component of stationary function $\phi_0$ of the differential part of diffusion operator

$$D^{\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+} \phi_0(r) = 0$$

which looks as follows

$$\phi_0(r) = \xi_1 + \xi_2 I^{\alpha}_{0,+} \frac{(1-r)^{\alpha-1}}{\Gamma(\alpha)\, p(r)} = \xi_1 + \xi_2 \psi(\alpha, 0, r). \tag{31}$$

The next function is the following integral

$$\begin{aligned}\varphi(r) = I^{\alpha}_{0,+} I^{\alpha}_{1,-} 1 = I^{\alpha}_{0,+} \frac{(1-r)^{\alpha}}{\Gamma(\alpha+1)} = (1-0)^{\alpha}(r-0)^{\alpha} \\ \times {}_1\Psi_2 \left[ \begin{array}{c|c} (1,1) & \\ (\alpha+1,-1) \quad (\alpha+1,1) & -\dfrac{r-0}{1-0} \end{array} \right].\end{aligned} \tag{32}$$

Again, using the Theorem 4 and calculating parameters according to (30)

$$\Delta = -1, \quad \delta = 1, \quad \mu = 2\alpha + \frac{1}{2}.$$

we conclude

$$\alpha > 0 \implies \mu > \frac{1}{2}$$

and the obtained Fox-Wright function (32) is continuous in interval $(0, 1]$ for any positive order $\alpha$.

**Lemma 12.** Let $\alpha > 1/2$ and denote

$$Y_\lambda(y) = \left( -\frac{2}{r} + \frac{l(l+1)}{r^2} + q_0(r) \right) y_\lambda(r) + \lambda w_\alpha y_\lambda(r)$$

$$\tilde{\Delta} = d_2 + d_1 \psi(\alpha, 0, 1).$$

Assume $\tilde{\Delta} \neq 0$. Then, on the $C[a,b]$ -space, the regular $(12-14)$ is equivalent to the integral equation

$$\begin{aligned} y_\lambda(r) = -I^{\alpha}_{0,+} \tfrac{1}{p(r)} I^{\alpha}_{1,-} Y_\lambda(y) \\ + A(r) \left( I^{\alpha}_{0,+} \tfrac{1}{p(r)} I^{\alpha}_{1,-} Y_\lambda(y) \right) |_{r=1} \end{aligned} \tag{33}$$

where the coefficient $A(r)$ is

$$A(r) = \frac{d_1}{\tilde{\Delta}} \psi(\alpha, 0, r)$$

and functions $\psi$ is defined in $(31)$.

**Proof.** By aid of composition rules, equation (12) can be rewriten as follows:

$$D^{\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+} \left[ y_\lambda(r) + I^{\alpha}_{0,+} \frac{1}{p(r)} I^{\alpha}_{1,-} Y_\lambda(y) \right] = 0.$$

The last equality refers that on the $C[a,b]$ -space the function in brackets is a stationary function of fractional Sturm Liouville problem for hydrogen atom operator. $D^{\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+}$ which according to (31) can be found as

$$\phi_0 = \xi_1 + \xi_2 I^{\alpha}_{0,+} \frac{(1-r)^{\alpha-1}}{\Gamma(\alpha)\, p(r)} = \xi_1 + \xi_2 \psi(\alpha, 0, r)$$

equation (12) in the form of

$$y_\lambda(r) + I^{\alpha}_{0,+} \frac{1}{p(r)} I^{\alpha}_{1,-} Y_\lambda(y) = \xi_1 + \xi_2 \psi(\alpha, 0, r), \tag{34}$$

to end the proof we should connect coefficients $\xi_j$, $j = 1, 2$ values $d_j$, $j = 1, 2$ determining the boundary conditions $(13 - 14)$. Let us note that the following formula results from composition rules (9) and equation (34)

$$I^{1-\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+} y_\lambda(r) = -I^{1}_{1,-} Y_\lambda(y) + \xi_2,$$

for continuous function $y_\lambda$, we obtain the following values a the ends

$$I^{1-\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+} y_\lambda(r) \Big|_{r=0} = -\int_0^1 Y_\lambda(y) + \xi_2 \tag{35}$$

$$I^{1-\alpha}_{1,-} p(r)^{C} D^{\alpha}_{0,+} y_\lambda(r) \Big|_{r=1} = \xi_2 \tag{36}$$

respectively for $y_\lambda$, using (34) we find

$$y_\lambda(0) = \phi_0(0) = \xi_1 \tag{37}$$

$$\begin{aligned} y_\lambda(1) &= \phi_0(1) - I^{\alpha}_{0,+} \frac{1}{p(r)} I^{\alpha}_{1,-} Y_\lambda(y) \Big|_{r=1} \\ &= \xi_1 + \xi_2 \psi(\alpha, 0, 1) - I^{\alpha}_{0,+} \frac{1}{p(r)} I^{\alpha}_{1,-} Y_\lambda(y) \Big|_{r=1}. \end{aligned} \tag{38}$$

The following set of linear equations for coefficients $\xi_j$ results from $(35 - 39)$

$$\begin{aligned} \xi_1 &= 0 \\ d_1 \xi_1 + \xi_2 (d_2 + d_1 \psi(\alpha, 0, 1)) &= d_1 F \end{aligned} \tag{39}$$

where $F = I^{\alpha}_{0,+} \frac{1}{p(r)} I^{\alpha}_{1,-} Y_\lambda(y) \Big|_{r=1}$.

Since $\tilde{\Delta} \neq 0$, the solution for coefficients $\xi_j$ $(j = 1, 2)$ is unique:

$$\xi_1 = 0$$

$$\xi_2 = d_1 F / \tilde{\Delta}$$

substituting the above solution into (34) we recover the equivalent integral equation $(33)$.

Furthermore we give notation such as

$$m_p = \min_{r \in (0,1]} |p(r)|,$$

$$A = \|A(r)\|, \quad M_\varphi = \|\varphi(r)\|,$$

where $\|.\|$ denotes the supremum norm on space $C[0,1]$ .$\zeta$

**Theorem 13.** Let $\alpha > 1/2$. Suppose that $\tilde{\Delta} \neq 0$. Then unique continuous eigenfunction $y_\lambda$ of singular fractional Sturm-Liouville problem for hydrogen atom operator $(12-14)$ corresponding to each eigenvalue obeying

$$\left\| \left( -\frac{2}{r} + \frac{l(l+1)}{r^2} + q_0(r) \right) + \lambda w_\alpha \right\| < \frac{m_p}{M_\varphi + B\varphi(1)} \quad (40)$$

exists and such eigenvalue is simple.

**Proof.** We have to say that equation (33) can be interpreted as a fixed point condition on function space $C[0,1]$,

$$y_\lambda = Ty_\lambda$$

where the mapping on the right-hand side is for any continuous function $g \in C[0,1]$ defined as

$$Tg = -I_{0,+}^{\alpha} \frac{1}{p(r)} I_{1,-}^{\alpha} Y_\lambda(y) + A(r) \left( I_{0,+}^{\alpha} \frac{1}{p(r)} I_{1,-}^{\alpha} Y_\lambda(y) \right) |_{r=1}$$

The following inequality will be useful in further estimations:

$$\begin{aligned} \|Y_\lambda(g) - Y_\lambda(\rho)\| &= \Bigg\| \Bigg( \left( -\frac{2}{r} + \frac{l(l+1)}{r^2} + q_0(r) \right) g \\ &+ \lambda w_\alpha g \Bigg) - \left( \left( -\frac{2}{r} + \frac{l(l+1)}{r^2} + q_0(r) \right) \rho + \lambda w_\alpha \rho \right) \Bigg\| \\ &= \left\| (g-\rho) \left( \left( -\frac{2}{r} + \frac{l(l+1)}{r^2} + q_0(r) \right) + \lambda w_\alpha \right) \right\| \\ &\le \|g-\rho\| \left\| \left( -\frac{2}{r} + \frac{l(l+1)}{r^2} + q_0(r) \right) + \lambda w_\alpha \right\|. \end{aligned}$$

By performing necassary operations for distance between images $T g$ and $T\rho$ for a pair of arbitrary continuous functions $g, \rho \in C[0,1]$.

$$\begin{aligned} \|Tg - T\rho\| &= \Bigg\| \Bigg[ -I_{0,+}^{\alpha} \frac{1}{p(r)} I_{1,-}^{\alpha} Y_\lambda(g) \\ &+ A(r) \left( I_{0,+}^{\alpha} \frac{1}{p(r)} I_{1,-}^{\alpha} Y_\lambda(g) \right)_{r=1} \Bigg] \\ &- \left[ -I_{0,+}^{\alpha} \frac{1}{p(r)} I_{1,-}^{\alpha} Y_\lambda(\rho) + A(r) \left( I_{0,+}^{\alpha} \frac{1}{p(r)} I_{1,-}^{\alpha} Y_\lambda(\rho) \right) |_{r=1} \right] \Bigg\| \\ &= \Bigg\| -I_{0,+}^{\alpha} \frac{1}{p(r)} I_{1,-}^{\alpha} \left( Y_\lambda(g) - Y_\lambda(\rho) \right) \\ &+ A(r) \left( I_{0,+}^{\alpha} \frac{1}{p(r)} I_{1,-}^{\alpha} \left( Y_\lambda(g) - Y_\lambda(\rho) \right) \right)_{r=1} \Bigg\| \\ &\le \|g-\rho\| \left\| \left( -\frac{2}{r} + \frac{l(l+1)}{r^2} + q_0(r) \right) + \lambda w_\alpha \right\| \left( \frac{M_\varphi}{m_p} + \frac{A\varphi(1)}{m_p} \right) \\ &\le \|g-\rho\| L \end{aligned}$$

where $L = \left\| \left( -\frac{2}{r} + \frac{l(l+1)}{r^2} + q_0(r) \right) + \lambda w_\alpha \right\| \left( \frac{M_\varphi}{m_p} + \frac{B\varphi(1)}{m_p} \right)$. By means of $(40)$, we state that mapping $T$ is a contraction on space $\langle C[0,1], \|.\| \rangle$

$$\|Tg - T\rho\| \le \|g-\rho\| L, \quad L \in (0,1).$$

Hence, a unique fixed point enounced as $y_\lambda \in C(0,1]$ exists that solves equation $(12, 33)$ and satisfies the boundary conditions $(13, 14)$, provided $(40)$ is applied. In that case such eigenvalues are simple. The proof is completed.

## Conclusion

In this article, we have extended the scope of some properties of fractional Sturm-Liouville problem for hydrogen atom operator. We pointed that its eigenvalues related to the fractional Sturm-Liouville problem for hydrogen atom operator. with the certain boundary conditions are real and its eigenfunctions corresponding to distinct eigenvalues are ortohogonal. Furthermore, we showed that the operator is self adjoint. Our new results are original and important for the fractional Sturm-Liouville theory.

## References

1. R., S. Johnson, An introduction to Sturm-Liouville theory. University of Newcastle (2006).
2. A., Zettl, Sturm-Liouville Theory, Mathematical Surveys and Monographs, Vol. 121. American Mathematical Society, (2005).
3. W., O. Amrein, A. M. Hinz, and D. B. Pearson (Editors), Sturm-Liouville Theory, Past and Present, Basel, Switzerland: Birkhauser, (2005).
4. E.S. Panakhov. R. Yilmazer, "A Hochstadt - Lieberman Theorem for the Hydrogen Atom Equation" App.and Comp. Math. Vol. 11, pp.74-80, (2012).
5. B., M. Levitan and I.S. Sargsjan, Introduction to Spectral Theory: Self adjoint Ordinary Differential Operators. American Math. Soc., Pro., R.I., (1975).
6. J. Qi., Sh. Chen. "Eigenvalue problems of the model from nonlocal continuum mechanics," J. Math. Phys. Vol. 52, No 073516, (2011).
7. A., Carpinteri and F. Mainardi (Editors), Fractals and Fractional Calculus in Continum Mechanics, Telos: Springer-Verlag, (1998).
8. B. J., West, M. Bologna, P. Grigolini Physics of Fractal Operators, New York, NY: Springer Verlag. (2003).
9. I., Podlubny, Fractional Differential Equations, San Diego, CA, USA: Academic Press, (1999).
10. R., Hilfer, (Ed.) Applications of Fractional Calculus in Physics, Singapore: World Scientific, (2000).

11. T. M., Atanackovic, B. Stankovic, "Generalized wave equation in nonlocal elasticity," Acta Mech. Vol. 208, pp. 1-10, (2009).

12. S. G., Samko, A. A. Kilbass, and O. I. Marichev, Fractional Integrals and Derivatives: Theory and Applications, PA: Gordon and Breach, (1993).

13. K. S., Miller and B. Ross, An Introduction to the Fractional Calculus and Fractional Differential Equations, New York, NY:John Wiley and Sons, Inc., (1993).

14. A. A., Kilbas, H. M. Srivastava, and J. J. Trujillo, Theory and Applications of Fractional Differential Equations, Amsterdam, Netherlands:Elsevier, (2006).

15. R., Yılmazer, E., Baş "Fractional Solutions of Confluent Hypergeometric Equation" J. Chung. Math. Soc. Vol. 25-2, (2012).

16. S. Das, R. Kumar, P.K. Gupta "Analytical Approximate Solution of Space-Time Fractional Diffusion Equation with a Moving Boundary Condition" Zeitschrift für Naturf. A Vol.66 281-289, (2011).

17. S. Kumar, H.Kocak, A.Yıldırım "The Solution of the Linear Fractional Partial Differential Equations Using the Homotopy Analysis Method" Zeitschrift für Naturf. A Vol. 67a 389-397 (2012).

18. M., Klimek, On Solutions of Linear Fractional Differential Equations of a Variational Type, Czestochowa: The Publishing Office of Czestochowa, University of Technology, (2009).

19. Q. M., Al-Mdallal, " An efficient method for solving fractional Sturm-Liouville problems," Chaos Solitons and Fractals, Vol. 40, pp. 183-189, (2009).

20. V. S., Erturk, "Computing eigenelements of Sturm-Liouville Problems of fractional order via fractional differential transform method," Mathematical and Computational Applications, Vol. 16, pp. 712-720, (2011).

21. M., Klimek and O. P., Argawal, , "On a Regular Fractional Sturm-Liouville Problem with Derivatives of Order in (0,1)" 13 th Int. Cont. Conf. (2012) .

22. A., Granas and J., Dugundji, Fixed Point Theory Springer-Verlag, New York, (2003).